



# STUDY OF THE RELATIONSHIP BETWEEN OLFACTORY PERCEPTION AND TOXICITY


## 1.- INTRODUCTION




## 2.- OBJECTIVES



## 3.- METHODOLOGY



## 4.- RESULTS AND CONCLUSIONS



### BIBLIOGRAPHY:

- Buck, L. & Axel, R. (1991). A novel multigene family may encode odorant receptors: a molecular basis for odor recognition. Cell, 65, 175-187.
- Cashdan, E. (1998). Adaptiveness of food learning and food aversions in children. Social Science Information, 37, 613-632.
- Haddad, R.; Weiss, T.; Khan, R.; Nadler, B.; Mandairon, N.; Bensafi, M.; Schneidman, E. & Sobel, N. (2010b). Global features of neural activity in the olfactory system form a parallel code that predicts olfactory behavior and perception. J. Neuroscience, 30 (27), 9017-9026.
- Zarzo, M. & Stanton, D. T. (2006). Identification of latent variables in a semantic odor profile database using principal component analysis. Chem. Senses, 31, 713-724.
- Zarzo, M. (2007). The sense of smell: molecular basis of odorant recognition. Biol. Rev., 82, 455-479.
- Zarzo, M. (2011). Hedonic judgments of chemical compounds are correlated with molecular size. Sensors, 11, 3667-3686.